



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

œsophageal ganglion, joins the miniature globe and expands into a retina. The author was unable to trace an opening through the pigment for the passage of light, but thinks it probable that, as in the ocelli of insects, such an aperture exists in the central part. The auditory capsules are situated at some distance behind the eyes, and may be distinctly seen with the microscope when the surrounding parts are carefully removed with fine needles. They are of a rounded or oval form, and each contains a beautifully transparent and highly refracting otolith, much larger than the lens of the eye.

The paper was accompanied with drawings illustrating the principal points mentioned in the description.

II. A paper was in part read, entitled, "On the Development of Striated Muscular Fibre in Mammalia." By WILLIAM S. SAVORY, M.D., F.R.C.S., Tutor of St. Bartholomew's Hospital Medical College. Communicated by JAMES PAGET, Esq. Received December 9, 1854.

The Society then adjourned over the Christmas recess, to meet again on the 11th of January next.

January 11, 1855.

THOMAS BELL, Esq., V.P., in the Chair.

I. The reading of Mr. SAVORY's paper "On the Development of Muscular Fibre in Mammalia," was resumed and concluded.

The author's observations were made chiefly upon fœtal pigs, but they have been confirmed by repeated examinations of the embryos of many other animals, and of the human fœtus.

If a portion of tissue immediately beneath the surface from the dorsal region of a fœtal pig, from one to two inches in length, be

examined microscopically, there will be seen, besides blood-corpuscles in various stages of development, nucleated cells and free nuclei or cytoblasts scattered through a clear and structureless blastema in great abundance. These cytoblasts vary in shape and size; the smaller ones, which are by far the most numerous, being generally round, and the larger ones more or less oval. Their outline is distinct and well defined, and one or two nucleoli may be seen in their interior as small, bright, highly refracting spots. The rest of their substance is either uniformly nebulous or faintly granular.

The first stage in the development of striated muscular fibre consists in the aggregation and adhesion of the cytoblasts, and their investment by blastema so as to form elongated masses. In these clusters the nuclei have, at first, no regular arrangement. Almost, if not quite as soon as the cytoblasts are thus aggregated, they become invested by the blastema, and this substance at the same time appears to be much condensed, so that many of the nuclei become obscured.

These nuclei, thus aggregated and invested, next assume a much more regular position. They fall into a single row with remarkable uniformity, and the surrounding substance at the same time grows clear and more transparent, and is arranged in the form of two bands bordering the fibre and bounding the extremities of the nuclei, so that now they become distinctly visible. They are oval, and form a single row in the centre of the fibre, closely packed together side by side, their long axes lying transversely, and their extremities bounded on either side by a thin clear pellucid border of apparently homogeneous substance.

It is to be observed how closely the muscular fibres of mammalia at this period of their development resemble their permanent form in many insects.

The fibres next increase in length and the nuclei separate. Small intervals appear between them. The spaces rapidly widen, until at last the nuclei lie at a very considerable distance apart. At the same time the fibre strikingly decreases in diameter; for as the nuclei separate, the lateral bands fall in and ultimately coalesce.

This lengthening of the fibre and consequent separation of the nuclei is due to an increase of material, and not to a stretching of the fibre.

Soon after the nuclei have separated some of them begin to decay.

They increase in size; their outline becomes indistinct; a bright border appears immediately within their margin; their contents become decidedly granular; their outline is broken and interrupted; and presently an irregular cluster of granules is all that remains, and these soon disappear.

It sometimes happens that the nuclei perish while in contact, before the fibre elongates; but the subsequent changes are the same.

The striæ generally first become visible at this period, immediately within the margin of the fibre.

The fibre is subsequently increased in size, and its development is continued by means of the surrounding cytoblasts. These attach themselves to its exterior, and then become invested by a layer of the surrounding blastema. Thus, as it were, nodes are formed at intervals on the surface of the fibre. These invested nuclei are at first readily detached, but they soon become intimately connected and indefinitely blended with the exterior of the fibre. All its characters are soon acquired, the nuclei at the same time gradually sink into its substance, and an ill-defined elevation, which soon disappears, is all that remains.

Lastly, the substance of the fibre becomes contracted and condensed. The diameter of a fibre towards, or at the close of intra-uterine life, is considerably less than at a much earlier period.

At the period of birth muscular fibres vary much in size.

The several stages in the development of muscular fibre, above mentioned, do not succeed each other as a simple consecutive series; on the contrary, two, or more, are generally progressing at the same time. Nor does each commence at the same period in all cases.

II. "On the General Integrals of the Equations of the Internal Equilibrium of an Elastic Solid." By WILLIAM JOHN MACQUORN RANKINE, Civil Engineer, F.R.SS. Lond. & Edinb., &c. Received December 7, 1854.

The *First Section* of this paper is introductory, containing a summary of principles already known respecting the elasticity of solids. Those principles are treated as the consequences of the following